

Trypanocidal Activity of Some Medicinal Plants Against the Epimastigotes of *Trypanosoma cruzi*

Gohari AR (Ph.D.)¹, Saeidnia S (Ph.D.)^{1*}, Hadjiakhoondi A (Ph.D.)^{1,2}, Naghinejad A (Ph.D.)³, Yagura T (Ph.D.)⁴

1- Medicinal Plants Research Center, Faculty of Pharmacy, Medical Sciences/ University of Tehran, Tehran, Iran

2- Department of Pharmacognosy, Faculty of Pharmacy, Medical Sciences/ University of Tehran, Tehran, Iran

3- Department of Botany, Faculty of Sciences, University of Tehran, Tehran, Iran

4- Department of Pharmacognosy, Graduate School of Pharmaceutical Sciences, Kyoto University, Kyoto, Japan

*Corresponding author: Medicinal Plants Research Center, Medical Sciences/ University of Tehran, P.O.Box: 14155-6451, Tehran, Iran, Tel: +98-21-66959090, Fax: +98-21-66461178

E-mail: soodabehsaeidnia@hotmail.com

Receive: 21 Jan. 2008

Acceptance: 17 Feb. 2008

Abstract

Background: Some of medicinal plants are a potential source of new drugs to improve the treatment of Chagase disease whose treatment is still a challenge. Here in this screening, the *in vitro* trypanocidal activity of some fractions for 16 medicinal plants, collected from the northern part of Iran, has been reported.

Methods: Aerial parts of the plants were dried carefully and followed by extraction with hexane and methanol, successively, by maceration at room temperature. Different concentrations of the plant extracts in ethanol were investigated against the epimastigotes of *T. cruzi*. The movement of epimastigotes was observed under a microscope. We assumed that immobilized organisms were died. The negative control contained ethanol in the same proportion utilized to dissolve the drugs. Each assay was performed in duplicate together with gentian violet as a positive control.

Results: Results show that hexane extracts of *Rubus hyrcanus* and *Salvia sclerae* have been observed the most activity against the epimastigotes of *T. cruzi* (MLC = 12.5 µg/ml). None of methanol fractions shows trypanocidal activity except *Salvia sclerae* (MLC = 50 µg/ml). *Echium amoenum*, *Satureja macrantha*, *S. atropatana* and *Stachys laxa* did not display activity lower than 100 µg/ml in both hexane and methanol extracts.

Conclusions: Some of Iranian medicinal plants (*Salvia sclerae*, *Marrubium vulgare* and *Rubus hyrcanus*) could be the promising source of active components against the epimastigotes of *T. cruzi* and need to further phytochemical and pharmacological studies.

Keywords: Trypanocidal activity, *Salvia sclerae*, *Rubus hyrcanus*



Introduction

Some of medicinal plants are a potential source of new drugs to improve the treatment of Chagase disease whose treatment is still a challenge and leading to approximately 400,000 deaths per year. Chagas disease is caused by the flagellate protozoan *Trypanosoma cruzi* (Trypanosomatina) [1]. Trypomastigotes ingested by the insect differentiates in to the proliferative epimastigote and then to metacyclic trypomastigotes. The latter form undergoes differentiation in to amastigotes, which after several reproductive cycles transform to trypomastigotes, the form responsible for the dissemination of the infection [2]. Current treatment is unsatisfactory, because the only two drugs are available, benznidazole and nifortimox, possess severe side effects and their activity is limited to the acute phase [3].

Literature reviews show that until recently, the *in vitro* trypanocidal activity of herbal extracts belonging to the families Meliaceae and Rutaceae has been evaluated. Among them, leaves of *Conchocarpus heterophyllus* and branches of *Trichilia ramalhoi* were the most active [1]. The extracts of the plants used in traditional Chinese medicine were tested *in vitro* with the epimastigote of *T. cruzi*, of which, *Lithospermum erythrorhizon*, *Saussurea lappa*, *Melia toosendan* and *Cinnamomum cassia* showed the greatest inhibitory activity of 100% [4]. Crude extracts (in the concentration 500 µg/ml) of *Bertholletia excelsa* stem barks were tested and showed significant *in vitro* trypanocidal activity against trypomastigote of *T. cruzi* [5]. Also, crude hydro alcoholic extracts and several fractions of 13 plants from Brazilian Atlantic Rain Forest were tested against epimastigote and trypomastigote forms of *T.*

cruzi and show a promising *in vitro* activity [6].

Previously, we reported the trypanocidal activity of some Iranian medicinal plants and isolated many active constituents of them. The most effective of those plants was *Dracocephalum kotschyi* and *D. subcapitatum* [7, 8, 9]. In this work, we assessed the effect of several extracts obtained from 16 plants, used in Iranian traditional medicine, against the epimastigotes of *T. cruzi*, the causative agent of chagas disease.

Experimental

Plant materials

All the plants were collected from northern part of Iran, including west Azerbyjan (near Uroomieh), Gilan (Ganjeh), Ghazvin (Telegan mountains), Golestan (National Park) and Mazandaran (Savad Kuh) provinces) in May and June (2004) during full flowering stage and identified by Dr. A. R. Naghinejad. The voucher herbarium specimens were deposited at the herbarium of Medicinal Plants Research Center housed in Faculty of Pharmacy, Tehran University of Medical Sciences.

Extractions of the plants

Aerial parts of the plants were dried and reduced to small pieces, followed by extraction with hexane and methanol, successively, by maceration (two time for each solvent and 24 h for each time) at room temperature. Then the solvents evaporated under reduced pressure to obtain the concentrated extracts and dried under vacuum in order to give dried powder of the extracts.

Evaluation of anti- epimastigote activity

Epimastigotes of *T. cruzi* (Tulahuen strain, prepared from Kyoto University, Japan) were

kept in GIT medium (Wako Pure Chemical Industrial, Ltd, Osaka, Japan) supplemented with hemin (12.4 μ M, Wako). The epimastigotes in GIT medium (10 μ L) were incubated with a test sample dissolved in EtOH (5 μ L) and autoclaved saline (185 μ L). Test samples contained different concentrations of the plant extracts prepared using serial dilution (2.5–200 μ g/ml). All samples were incubated at 27 °C for 24 h. The movement of epimastigotes was observed under a microscope. We assumed that immobilized organisms were dead. The control contained ethanol in the same proportion as used to dissolve the extracts. Each assay was performed in duplicate. Minimum lethal

concentration (MLC, concentration on which, all epimastigotes were dead) of each fraction was performed. Gentian violet (MLC = 6.3 μ M) was used as a positive control [8, 9].

Results and Discussion

The results of *in vitro* trypanocidal activity of some Iranian medicinal plants are summarized in table 1. In this study, we prepared two different extracts (hexane and methanol) of each plant except two species of *Achillea*. Because, the hexane extract of those mentioned plants was too waxy and not dissolved in safely amounts of DMSO (below 1%) for epimastigotes as a co-solvent, we tried to prepare ethyl acetate extract.

Table 1. Comparative results of trypanocidal activity from different extracts of the plants against the epimastigotes of *T. cruzi*.

NO	Plant name	Family	Hexane MLC (μ g/ml)	MeOH MLC (μ g/ml)	AcOEt MLC (μ g/ml)
1	<i>Achillea talagonica</i>	Asteraceae	-	200	50
2	<i>Achillea tenuifolia</i>	Asteraceae	-	200	50
3	<i>Tanacetum polycephalum</i>	Asteraceae	50	200	-
4	<i>Rubus hyrcanus</i>	Rosaceae	12.5	>200	-
5	<i>Rubus discolor</i>	Rosaceae	50	200	-
6	<i>Onosma bulbosum</i>	Boraginaceae	50	>200	-
7	<i>Echium amoenum</i>	Boraginaceae	200	100	-
8	<i>Marrubium vulgare</i>	Lamiaceae	25	100	-
9	<i>Stachys laxa</i>	Lamiaceae	100	100	-
10	<i>Thymus kotschyianus</i>	Lamiaceae	50	100	-
11	<i>Leonurus cardiaca</i>	Lamiaceae	50	100	-
12	<i>Salvia sclarea</i>	Lamiaceae	12.5	50	-
13	<i>Satureja mutica</i>	Lamiaceae	50	100	-
14	<i>Satureja atrapatana</i>	Lamiaceae	100	100	-
15	<i>Satureja macrantha</i>	Lamiaceae	100	200	-
16	<i>Satureja spicigera</i>	Lamiaceae	50	100	-

Results show that none of methanol fractions displayed trypanocidal activity except *Salvia sclerae*. Hexane extracts of *Rubus hyrcanus* and *Salvia sclerae* have been observed the most activity against the epimastigotes of *T. cruzi* (MLC = 12.5 µg/ml). Hexane fraction of *Marrubium vulgare* was active with MLC = 25 µg/ml. Hexane extracts of *Tanacetum polycephalum*, *Rubus discolor*, *Onosma bulbotrichum*, *Thymus kotschyianus*, *Satureja mutica* and *S. spicigera* showed effects with minimum lethal concentration at 50 µg/ml. Ethyl acetate fractions of *Achillea* species were effective as well. Among the 16 plants, *Stachys laxa*, *Echium amoenum*, *Satureja macrantha* and *S. atropatana* did not represent activity in any fraction tested.

There are a few reports about the presence of pentacyclic triterpenes like oleanolic acid and ursolic acid in the *Rubus chingii* and *Salvia sclarea* extracts [10, 11]. Also there are some reports representing trypanocidal activity of ursan skeleton [12, 13], so it seems that trypanocidal activity of hexane extracts of *Salvia sclarea* and *Rubus hyrcanus* is some part due to the presence of these triterpene acid components.

Acknowledgements

The authors wish to thank Prof. Gisho Honda and Dr. Michiho Ito from Kyoto University for cooperating in this project and Research council of Tehran University of Medical Sciences.

References

1. Ambrozin ARP, Vieira PC, Fernandes JB, Fernandes da silva MFG and de Albuquerque S. Trypanocidal activity of Meliaceae and Rutaceae plant extracts. *Mem. Inst. Oswaldo cruz.* 2004; 99: 227-31.
2. Coura JR and de Castro SL. A critical review on Chagas disease chemotherapy. *Mem. Inst. Oswaldo cruz.* 2002; 97: 3 - 24.
3. Nogueira- Torres B, Rodriguez- Paez L, Ramirez IB and Ramirez CW. Trypanocidal activity of isopropyl salicylaldehyde and 4-isopropyl salicylic acid on *Trypanosoma cruzi*. *Rev. Lat. Microbiol.* 2001; 43: 1 - 6.
4. Lirussi D, Li J, Prieto JM, Gennari M, Buschiazzi H, Rios JL and Zaidenberg A. Inhibition of *Trypanosoma cruzi* by plant extracts used in Chinese medicine. *Fitoter.* 2004; 75: 718 - 23.
5. Campos FR, Januario AH, Rosas LV, Nascimento SK, Pereira PS, Franca SC, Cordeiro MS, Toldo MP and Albuquerque S. Trypanocidal activity of extracts and fractions of *Bertholletia excelsa*. *Fitoter.* 2005; 76: 26-9.
6. Pizzolatti MG, Koga AH, Grisard EC and Steindel M. Trypanocidal activity of extracts from Brazilian Atlantic Rain Forest plant species. *Phytomed.* 2003; 10: 422 - 6.
7. Gohari AR, Saeidnia S, Matsuo K, Uchiyama N, Yagura T, Ito M, Kiuchi F and Honda G. Flavonoid constituents of *Dracocephalum kotschy* growing in Iran and their trypanocidal activity. *J. Nat. Med.* 2003; 57: 250 - 2.
8. Saeidnia S, Gohari AR, Ito M, Kiuchi F and Honda G. Bioactive constituents from

Dracocephalum subcapitatum (O. Kunntze) Lipsky. *Z. Naturforsch.* 2005; 60: 22 - 4.

9. Saeidnia S, Gohari AR, Kiuchi F and Honda G. Anti-epimastigote activity of some Iranian medicinal plants. *Iran J. Pharmaceut. Res.* 2005; 2: 101 - 3.

10. Guo QL and Yang JS. Studies on chemical constituents in fruit of *Rubus chingii*. *Zhongguo Zhong Yao. Za. Zhi.* 2005; 30: 198 - 200.

11. Kuzma L, Skrzypek Z and Wysokinska H. Diterpenoids and triterpenoids in hairy roots of

Salvia sclarea. *Plant Cell, Tissue and Organ Culture*, 2006; 84: 100152 - 60.

12. Abe F, Yamauchi T, Nagao T, Kinjo J, Okabe H, Higo H and Akahane H. Ursolic acid as a trypanocidal constituents in rosemary. *Biol. Pharm. Bull.* 2002; 25: 1485- 7.

13. Saeidnia S, Gohari AR, Uchiyama N, Ito M, Honda G and Kiuchi F. Two new monoterpene glycosides and other trypanocidal terpenoids from *Dracocephalum kotschyi*. *Chem. Pharm. Bull.* 2004; 52: 1249 - 50.